

# Pipecolate, specific biomarker of lysine deficiency

Gaëtan Roisné-Hamelin<sup>a</sup>, Claire Gaudichon<sup>a</sup>, Sarita Devi<sup>b</sup>, Jean-Charles Martin<sup>c</sup>, Catherine Tardivel<sup>c</sup>, Anura Kurpad<sup>b</sup>, Daniel Tomé<sup>a</sup>, Delphine Jouan-Rimbaud Bouveresse<sup>a</sup> and Dalila Azzout-Marniche<sup>a</sup>

<sup>a</sup> Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, 91120, Palaiseau, France.


<sup>b</sup> Division of Nutrition, St John's Reserach Institute, St John's National academy of health sciences, bangalore, India.

<sup>c</sup> Aix Marseille université, INSERM, C2VN, Marseille, France.

## Background & Objective

- The consumption of poor-quality protein increases the risk of essential amino acid (EAA) deficiency, particularly for lysine, threonine and methionine. Thus, it is necessary to be able to detect easily EAA deficiency.
- We have previously identified pipecolate and taurine as potential biomarkers for lysine and threonine deficiency, respectively (Moro et al. 2023, J nutr, 153:2571-2584).
- The purpose of this study was to develop metabolomic approaches to identify specific biomarkers for an EAA deficiency.


## Methods





- Body weight and food intake (FI) were measured daily.
- 24h-urine was analyzed LC-MS metabolomic.
- Body composition was determined by EchoMRI.

- Body weight was analyzed by repeated measures, mixt model and other variables by one-way ANOVA.
- Metabolic features were analyzed by PLS-DA and individually test for diet effects by ANOVA.

## Results



Our results confirmed that protein and EAA deficiency induced growth retardation and supplementation permits to restore growth, but a delay of length, lean body mass and specific organs weights remains after supplementation.



Urinary metabolites from lysine degradation pathway, particularly pipecolate and N-N-N-Trimethyllysine, signed lysine deficiency.

- The urinary metabolome allowed to discriminate between the deficient and no deficient diet, and we were able to identify specific signatures for methionine and lysine deficiency.
- The best model retained 4 latent variables with LV3 allowing the discrimination of methionine deficient diets and LV4 allowing the discrimination of lysine deficient diets.
- Further analyses are required to investigate the specific signature for threonine intake.

## Conclusion

- Our results showed that **EAA deficiencies influence the urinary metabolome**.
- We identified **specific urinary metabolomic for lysine and methionine deficiency**.
- We confirmed that **pipecolate is a urinary biomarker that specifically signs lysine deficiency**.
- Our results showed that **the deficiency/supplementation method could be applied to identify specific EAA biomarkers**.
- The urinary biomarkers identified could be easily applied to detect EAA or protein status.

## Contact

- [dalila.azzout-marniche@agroparistech.fr](mailto:dalila.azzout-marniche@agroparistech.fr)
- [gaetan.hamelin@agroparistech.fr](mailto:gaetan.hamelin@agroparistech.fr)
- [delphine.bouveresse@agroparistech.fr](mailto:delphine.bouveresse@agroparistech.fr)